


# **Lecture 2. Diffusion models**

## **Introduction to Bayesian statistical Learning II**

**21.05.2025 Instructors: Alina Bazarova, Jose Robledo**

# Denoising Diffusion Probabilistic Models (DDPM)


Another type of generative models. What do they have to do with Bayes?



1. Forward diffusion process: gradually **adding noise** to samples

# Denoising Diffusion Probabilistic Models (DDPM)


Another type of generative models. What do they have to do with Bayes?



1. Forward diffusion process: gradually **adding noise** to samples
2. Reverse diffusion process: **recreating** the sample **from noise**

# Denoising Diffusion Probabilistic Models (DDPM)

Another type of generative models. What do they have to do with Bayes?



1. Forward diffusion process: gradually **adding noise** to samples
2. Reverse diffusion process: **recreating** the sample **from noise**
3. Heavily relying on conditional probability and **Bayes theorem** in particular

## Forward diffusion process

Let  $x_0 \sim q(x)$  be a real data distribution.

We produce a sequence of **noisy samples**  $x_1, \dots, x_T$

# Forward diffusion process

Let  $x_0 \sim q(x)$  be a real data distribution.

We produce a sequence of **noisy samples**  $x_1, \dots, x_T$

Noise steps controlled by a **variance schedule**  $\{\beta_t \in (0,1)\}_{t=1}^T$

## Forward diffusion process

Let  $x_0 \sim q(x)$  be a real data distribution.

We produce a sequence of **noisy samples**  $x_1, \dots, x_T$

Noise steps controlled by a **variance schedule**  $\{\beta_t \in (0,1)\}_{t=1}^T$

$$q(x_t | x_{t-1}) \sim \mathcal{N}(x_t; \sqrt{1 - \beta_t} x_{t-1}, \beta_t \mathbf{I}), \quad q(x_{1:T} | x_0) = \prod_{t=1}^T q(x_t | x_{t-1})$$

## Forward diffusion process

Let  $x_0 \sim q(x)$  be a real data distribution.

We produce a sequence of **noisy samples**  $x_1, \dots, x_T$

Noise steps controlled by a **variance schedule**  $\{\beta_t \in (0,1)\}_{t=1}^T$

$$q(x_t | x_{t-1}) \sim \mathcal{N}(x_t; \sqrt{1 - \beta_t} x_{t-1}, \beta_t \mathbf{I}), \quad q(x_{1:T} | x_0) = \prod_{t=1}^T q(x_t | x_{t-1})$$

As  $T \rightarrow \infty$ ,  $x_T$  is equivalent to a **Gaussian distribution**



## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

Let  $\alpha_t = 1 - \beta_t$  and  $\bar{\alpha}_t = \prod_{i=1}^t \alpha_i$

## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

Let  $\alpha_t = 1 - \beta_t$  and  $\bar{\alpha}_t = \prod_{i=1}^t \alpha_i$

$$x_t = \sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \sqrt{\alpha_t\alpha_{t-1}}x_{t-2} + \sqrt{1 - \alpha_t\alpha_{t-1}}\bar{\varepsilon}_{t-2}$$

## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\bar{\alpha}_t} x_{t-1} + \sqrt{1 - \alpha_t} \varepsilon_{t-1} = \sqrt{\alpha_t \bar{\alpha}_{t-1}} x_{t-2} + \sqrt{1 - \alpha_t \bar{\alpha}_{t-1}} \bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t} x_{t-1} + \sqrt{1 - \alpha_t} \varepsilon_{t-1} = \sqrt{\alpha_t \alpha_{t-1}} x_{t-2} + \sqrt{1 - \alpha_t \alpha_{t-1}} \bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

$$\sqrt{\alpha_{t-1}} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2} \sim \mathcal{N}(0, \alpha_{t-1} - \alpha_t \alpha_{t-1} + 1 - \alpha_{t-1})$$

## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t} x_{t-1} + \sqrt{1 - \alpha_t} \varepsilon_{t-1} = \sqrt{\alpha_t \alpha_{t-1}} x_{t-2} + \sqrt{1 - \alpha_t \alpha_{t-1}} \bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

$$\sqrt{\alpha_{t-1}} \sqrt{1 - \alpha_t} \varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}} \varepsilon_{t-2} \sim \mathcal{N}(0, \cancel{\alpha_{t-1}} - \alpha_t \alpha_{t-1} + 1 - \cancel{\alpha_{t-1}})$$

## Forward diffusion process

Can we sample  $x_t$  at any arbitrary time step  $t$  ? Yes!!!

The key: good old **reparametrisation trick!**

$$\text{Let } \alpha_t = 1 - \beta_t \text{ and } \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

$$x_t = \sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \sqrt{\alpha_t\alpha_{t-1}}x_{t-2} + \sqrt{1 - \alpha_t\alpha_{t-1}}\bar{\varepsilon}_{t-2}$$

$$\text{NB: } \sqrt{\alpha_t}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2}, \quad \varepsilon_i \sim \mathcal{N}(0, \mathbf{I}), \text{ hence}$$

$$\sqrt{\alpha_{t-1}}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2} \sim \mathcal{N}(0, \alpha_{t-1} - \alpha_t\alpha_{t-1} + 1 - \alpha_{t-1})$$

$$\sqrt{\alpha_{t-1}}\sqrt{1 - \alpha_t}\varepsilon_{t-1} + \sqrt{1 - \alpha_{t-1}}\varepsilon_{t-2} \sim \sqrt{1 - \alpha_t\alpha_{t-1}}\mathcal{N}(0, \mathbf{I})$$

## Forward diffusion process

Following the same argument

$$x_t = \sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \dots = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I}), \quad \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$



## Forward diffusion process

Following the same argument

$$x_t\sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \dots = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I}), \quad \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

Hence  $q(x_t | x_0) \sim \mathcal{N}(x_t; \sqrt{\bar{\alpha}_t}x_0, (1 - \bar{\alpha}_t)\mathbf{I})$

## Forward diffusion process

Following the same argument

$$x_t\sqrt{\alpha_t}x_{t-1} + \sqrt{1 - \alpha_t}\varepsilon_{t-1} = \dots = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I}), \quad \bar{\alpha}_t = \prod_{i=1}^t \alpha_i$$

Hence  $q(x_t | x_0) \sim \mathcal{N}(x_t; \sqrt{\bar{\alpha}_t}x_0, (1 - \bar{\alpha}_t)\mathbf{I})$

Larger update step when the sample gets noisier:  $\beta_1 < \beta_2 < \dots < \beta_T$ ,

and therefore  $\bar{\alpha}_1 > \dots > \bar{\alpha}_T$

## Reverse diffusion process

If we reverse the above process and sample from  $q(x_{t-1} | x_t)$ ,

we can recreate a sample from  $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate  $q(x_{t-1} | x_t)$ . We do it with another probability density function  $p_\theta$

## Reverse diffusion process

If we reverse the above process and sample from  $q(x_{t-1} | x_t)$ ,

we can recreate a sample from  $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate  $q(x_{t-1} | x_t)$ . We do it with another probability density function  $p_\theta$

$$p_\theta(x_{0:T}) = p(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t),$$

## Reverse diffusion process

If we reverse the above process and sample from  $q(x_{t-1} | x_t)$ ,

we can recreate a sample from  $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate  $q(x_{t-1} | x_t)$ . We do it with another probability density function  $p_\theta$

$$p_\theta(x_{0:T}) = p(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t), \quad p_\theta(x_{t-1} | x_t) = \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$$

Which is learned via a neural network ( $\theta$  are the parameters of NN)!

## Reverse diffusion process

If we reverse the above process and sample from  $q(x_{t-1} | x_t)$ ,

we can recreate a sample from  $x_T \sim \mathcal{N}(0, \mathbf{I})$

We need to estimate  $q(x_{t-1} | x_t)$ . We do it with another probability density function  $p_\theta$

$$p_\theta(x_{0:T}) = p(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t), \quad p_\theta(x_{t-1} | x_t) = \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$$

Which is learned via a neural network ( $\theta$  are the parameters of NN)!

**Important:**  $q(x_{t-1} | x_t, x_0) \sim \mathcal{N}(x_{t-1}; \tilde{\mu}(x_t, x_0), \tilde{\beta}\mathbf{I})$  **tractable!**

**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$



**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$q(x_{t-1} | x_t, x_0) = q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right)$$

**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$\begin{aligned} q(x_{t-1} | x_t, x_0) &= q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right) \\ &= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right) \end{aligned}$$

**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$\begin{aligned} q(x_{t-1} | x_t, x_0) &= q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right) \\ &= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right) \end{aligned}$$

$$\text{Hence } \tilde{\beta}_t = 1/\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right) = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t,$$

**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$\begin{aligned} q(x_{t-1} | x_t, x_0) &= q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right) \\ &= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right) \end{aligned}$$

$$\text{Hence } \tilde{\beta}_t = 1/\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right) = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t} \beta_t,$$

$$\tilde{\mu}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)/\tilde{\beta}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)\frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t$$

**Conditioning on  $x_0$ . Bayes rule to the rescue!**  $P(A | B) = \frac{P(B)P(B | A)}{P(A)}$

$$P(A | B, C) = \frac{P(AB | C)}{P(B | C)} = \frac{P(B | A, C)P(A | C)}{P(B | C)}$$

$$q(x_{t-1} | x_t, x_0) = q(x_t | x_{t-1}, x_0) \frac{q(x_{t-1} | x_0)}{q(x_t | x_0)} \propto \exp\left(-\frac{1}{2}\left(\frac{(x_t - \sqrt{\alpha_t}x_{t-1})^2}{1 - \alpha_t} + \frac{(x_{t-1} - \sqrt{\bar{\alpha}_{t-1}}x_0)^2}{1 - \bar{\alpha}_{t-1}} - \frac{(x_t - \sqrt{\bar{\alpha}_t}x_0)^2}{1 - \bar{\alpha}_t}\right)\right)$$

$$= \exp\left(-\frac{1}{2}\left(\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right)x_{t-1}^2 - 2\left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)x_{t-1} + C(x_0, x_1)\right)\right)$$

$$\text{Hence } \tilde{\beta}_t = 1/\left(\frac{\alpha_t}{\beta_t} + \frac{1}{1 - \bar{\alpha}_{t-1}}\right) = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t,$$

$$\tilde{\mu}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)/\tilde{\beta}_t = \left(\frac{\sqrt{\alpha_t}}{\beta_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}}{1 - \bar{\alpha}_{t-1}}x_0\right)\frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}\beta_t$$

**We can reparametrise it further!  $\rightarrow$**

$$= \frac{\sqrt{\alpha_t}(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t}x_t + \frac{\sqrt{\bar{\alpha}_{t-1}}\beta_t}{1 - \bar{\alpha}_t}x_0$$

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$        $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$        $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$        $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t$



**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$        $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t$

**How are we going to do that?**

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$        $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t$

**How are we going to do that? Minimise negative log-likelihood  $-\log p_{\theta}(x_0)$**

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon,$        $x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t$

**How are we going to do that? Minimise negative log-likelihood  $-\log p_{\theta}(x_0)$**

$$-\log p_{\theta}(x_0) \leq -\log p_{\theta}(x_0) + D_{KL}(q(x_{1:T} | x_0) || p_{\theta}(x_{1:T} | x_0))$$

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_\theta(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$

So we will train  $\mu_\theta$  to predict  $\tilde{\mu}_t$

**How are we going to do that? Minimise negative log-likelihood  $-\log p_\theta(x_0)$**

**Kullback-Leibler divergence  $\geq 0$**

$-\log p_\theta(x_0) \leq -\log p_\theta(x_0) + D_{KL}(q(x_{1:T} | x_0) || p_\theta(x_{1:T} | x_0))$

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_\theta(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_\theta(x_t, t), \Sigma_\theta(x_t, t))$

So we will train  $\mu_\theta$  to predict  $\tilde{\mu}_t$

**How are we going to do that? Minimise negative log-likelihood  $-\log p_\theta(x_0)$**

**Kullback-Leibler divergence  $\geq 0$**

$-\log p_\theta(x_0) \leq -\log p_\theta(x_0) + D_{KL}(q(x_{1:T} | x_0) || p_\theta(x_{1:T} | x_0)) =$

$-\log p_\theta(x_0) + E_{x_{1:T} \sim q(x_{1:T} | x_0)}[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})/p_\theta(x_0)}] = -\log p_\theta(x_0) + E_q[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] + \log p_\theta(x_0)$

**Remember!**  $x_t = \sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon, \quad x_0 = \frac{1}{\sqrt{\bar{\alpha}_t}}(x_t - \sqrt{1 - \bar{\alpha}_t}\varepsilon_t)$

Substitute this into expression of  $\tilde{\mu}_t$  and get  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**Recap!** We need to learn the distributions  $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

So we will train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t$

**How are we going to do that? Minimise negative log-likelihood  $-\log p_{\theta}(x_0)$**

**Kullback-Leibler divergence  $\geq 0$**



$$-\log p_{\theta}(x_0) \leq -\log p_{\theta}(x_0) + D_{KL}(q(x_{1:T}|x_0) || p_{\theta}(x_{1:T}|x_0)) =$$

$$-\log p_{\theta}(x_0) + E_{x_{1:T} \sim q(x_{1:T}|x_0)}[\log \frac{q(x_{1:T}|x_0)}{p_{\theta}(x_{0:T})/p_{\theta}(x_0)}] = -\cancel{\log p_{\theta}(x_0)} + E_q[\log \frac{q(x_{1:T}|x_0)}{p_{\theta}(x_{0:T})}] + \cancel{\log p_{\theta}(x_0)} = E_q[\log \frac{q(x_{1:T}|x_0)}{p_{\theta}(x_{0:T})}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)} \log p_\theta(x_0)$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)} \log p_\theta(x_0)$$

We used that  $E_{q(x_0)}E_{q(x_{1:T}|x_0)}f(x_{0:T}) = E_{q(x_{0:T})}f(x_{0:T})$  for any  $f$  of our interest



$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)}p_\theta(x_0)$$

We used that  $E_{q(x_0)}E_{q(x_{1:T}|x_0)}f(x_{0:T}) = E_{q(x_{0:T})}f(x_{0:T})$  for any  $f$  of our interest

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)}p_\theta(x_0)$$

We used that  $E_{q(x_0)}E_{q(x_{1:T}|x_0)}f(x_{0:T}) = E_{q(x_{0:T})}f(x_{0:T})$  for any  $f$  of our interest

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})} [\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] \geq - E_{q(x_0)} p_\theta(x_0)$$

We used that  $E_{q(x_0)} E_{q(x_{1:T} | x_0)} f(x_{0:T}) = E_{q(x_{0:T})} f(x_{0:T})$  for any  $f$  of our interest

$$L_{VLB} = E_{q(x_{0:T})} [\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] = E_q [\log \frac{\prod_{t=1}^T q(x_t | x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t)}] = E_q [-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t | x_{t-1})}{p_\theta(x_{t-1} | x_t)}]$$

$$\text{Let } L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] \geq -E_{q(x_0)} p_\theta(x_0)$$

We used that  $E_{q(x_0)} E_{q(x_{1:T} | x_0)} f(x_{0:T}) = E_{q(x_{0:T})} f(x_{0:T})$  for any  $f$  of our interest

$$\begin{aligned} L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T} | x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t | x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1} | x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t | x_{t-1})}{p_\theta(x_{t-1} | x_t)}] \\ &= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t | x_{t-1})}{p_\theta(x_{t-1} | x_t)} + \log \frac{q(x_1 | x_0)}{p_\theta(x_0 | x_1)}] \end{aligned}$$

$$L_{VLB} = E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}]$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$



$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \log \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(x_1|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \log \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(\cancel{x_1}|x_0)} + \log \frac{q(\cancel{x_1}|x_0)}{p_\theta(x_0|x_1)}]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \log \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(\cancel{x_1}|x_0)} + \log \frac{q(\cancel{x_1}|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[\log \frac{q(x_T|x_0)}{p_\theta(x_T)} + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} - \log p_\theta(x_0|x_1)]
\end{aligned}$$

$$\begin{aligned}
L_{VLB} &= E_{q(x_{0:T})}[\log \frac{q(x_{1:T}|x_0)}{p_\theta(x_{0:T})}] = E_q[\log \frac{\prod_{t=1}^T q(x_t|x_{t-1})}{p_\theta(x_T) \prod_{t=1}^T p_\theta(x_{t-1}|x_t)}] = E_q[-\log p_\theta(x_T) + \sum_{t=1}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_t|x_{t-1})}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] = E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \dots] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \sum_{t=2}^T \frac{q(x_t|x_0)}{q(x_{t-1}|x_0)} + \log \frac{q(x_1|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[-\log p_\theta(x_T) + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} + \log \frac{q(x_T|x_0)}{q(\cancel{x_1}|x_0)} + \log \frac{q(\cancel{x_1}|x_0)}{p_\theta(x_0|x_1)}] \\
&= E_q[\log \frac{q(x_T|x_0)}{p_\theta(x_T)} + \sum_{t=2}^T \log \frac{q(x_{t-1}|x_t, x_0)}{p_\theta(x_{t-1}|x_t)} - \log p_\theta(x_0|x_1)] \\
&= E_q[\underbrace{D_{KL}(q(x_T|x_0) || p_\theta(x_T))}_{L_T} + \sum_{t=2}^T \underbrace{D_{KL}(q(x_{t-1}|x_t, x_0) || p_\theta(x_{t-1}|x_t))}_{L_{t-1}} - \underbrace{\log p_\theta(x_0|x_1)}_{L_0}]
\end{aligned}$$

## Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

## Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T|x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

## Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T | x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

$$L_t = D_{KL}(q(x_{t-1} | x_t, x_0) || p_{\theta}(x_{t-1} | x_t)) \quad \text{The main thing!}$$

## Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T|x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

$$L_t = D_{KL}(q(x_{t-1}|x_t, x_0) || p_{\theta}(x_{t-1}|x_t)) \quad \text{The main thing!}$$

$$L_0 = -\log p_{\theta}(x_0|x_1) \quad \text{Can be omitted or modelled via a separate decoder, derived from}$$

$$\mathcal{N}(x_0; \mu_{\theta}(x_1, 1), \Sigma_{\theta}(x_1, 1))$$



## Simplifying the loss further

$$L_{VLB} = L_T + L_{T-1} + \dots + L_0$$

$$L_T = D_{KL}(q(x_T | x_0) || p_{\theta}(x_T)) \quad \text{Constant, since no trainable parameters and } x_T \text{ is Gaussian noise}$$

$$L_t = D_{KL}(q(x_{t-1} | x_t, x_0) || p_{\theta}(x_{t-1} | x_t)) \quad \text{The main thing!}$$

$$L_0 = -\log p_{\theta}(x_0 | x_1) \quad \text{Can be omitted or modelled via a separate decoder, derived from}$$

$$\mathcal{N}(x_0; \mu_{\theta}(x_1, 1), \Sigma_{\theta}(x_1, 1))$$

**Remember again!** We need to learn the distributions  $p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$

That is for  $L_t$  !

## Parametrisation of $L_t$ for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

## Parametrisation of $L_t$ for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_t)$

**But  $x_t$  is available during training time!**

## Parametrisation of $L_t$ for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_t)$

**But  $x_t$  is available during training time!**

Hence, let us **predict the noise term instead!**

## Parametrisation of $L_t$ for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t)$

**But  $x_t$  is available during training time!**

Hence, let us **predict the noise term instead!**

$$\mu_{\theta}(x_t, t) = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_{\theta}(x_t, t))$$

## Parametrisation of $L_t$ for Training Loss

$$p_{\theta}(x_{t-1} | x_t) \sim \mathcal{N}(x_{t-1}; \mu_{\theta}(x_t, t), \Sigma_{\theta}(x_t, t))$$

We would like to train  $\mu_{\theta}$  to predict  $\tilde{\mu}_t = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_t)$

**But  $x_t$  is available during training time!**

Hence, let us **predict the noise term instead!**

$$\mu_{\theta}(x_t, t) = \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_{\theta}(x_t, t)) \quad x_{t-1} \sim \mathcal{N}(x_{t-1}; \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\epsilon_{\theta}(x_t, t)), \Sigma_{\theta}(x_t, t))$$

## Parametrisation of $L_t$ for Training Loss

$$L_t = E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_\theta(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_\theta(x_t, t)||^2 \right]$$

## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t)) ||^2 \right] \end{aligned}$$



## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}||_2^2} || \cancel{\frac{1}{\sqrt{\alpha_t}} x_t} - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t - \cancel{\frac{1}{\sqrt{\alpha_t}} x_t} - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2 \alpha_t (1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(x_t, t)||^2 \right] \end{aligned}$$

## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_{\theta}(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

**In practice can simplify even further!**

## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_\theta(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_\theta(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_\theta||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t)) ||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_\theta||_2^2} ||\varepsilon_t - \varepsilon_\theta(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_\theta||_2^2} ||\varepsilon_t - \varepsilon_\theta(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

**In practice can simplify even further!**

$$L_t^{simple} = E_{t, x_0, \varepsilon} [ ||\varepsilon_t - \varepsilon_\theta(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 ]$$

## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned} L_t &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}||_2^2} ||\frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_t) - \frac{1}{\sqrt{\alpha_t}}(x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}}\varepsilon_{\theta}(x_t, t))||^2 \right] \\ &= E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(x_t, t)||^2 \right] = E_{x_0, \varepsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 \right] \end{aligned}$$

**In practice can simplify even further!**

$$L_t^{simple} = E_{t, x_0, \varepsilon} [ ||\varepsilon_t - \varepsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\varepsilon_t, t)||^2 ]$$


**Bottom line: what we are doing is predicting the noise!**

## Parametrisation of $L_t$ for Training Loss

$$\begin{aligned}
 L_t &= E_{x_0, \epsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}(x_t, t)||_2^2} ||\tilde{\mu}_t(x_t, x_0) - \mu_{\theta}(x_t, t)||^2 \right] \\
 &= E_{x_0, \epsilon} \left[ \frac{1}{2 ||\Sigma_{\theta}||_2^2} || \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \epsilon_t) - \frac{1}{\sqrt{\alpha_t}} (x_t - \frac{1 - \alpha_t}{\sqrt{1 - \bar{\alpha}_t}} \epsilon_{\theta}(x_t, t)) ||^2 \right] \\
 &= E_{x_0, \epsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\epsilon_t - \epsilon_{\theta}(x_t, t)||^2 \right] = E_{x_0, \epsilon} \left[ \frac{(1 - \alpha_t)^2}{2\alpha_t(1 - \bar{\alpha}_t) ||\Sigma_{\theta}||_2^2} ||\epsilon_t - \epsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\epsilon_t, t)||^2 \right]
 \end{aligned}$$

In practice can simplify even further!

$$L_t^{simple} = E_{t, x_0, \epsilon} [ ||\epsilon_t - \epsilon_{\theta}(\sqrt{\bar{\alpha}_t}x_0 + \sqrt{1 - \bar{\alpha}_t}\epsilon_t, t)||^2 ]$$



**Bottom line: what we are doing is predicting the noise!** Unet architecture is used for that

## Parametrization of $\beta_t$

Typically a sequence of linearly increasing constants: e.g. from  $\beta_1 = 10^{-4}$  to  $\beta_T = 0.02$

## Parametrization of $\beta_t$

Typically a sequence of linearly increasing constants: e.g. from  $\beta_1 = 10^{-4}$  to  $\beta_T = 0.02$

Can be fixed (not learnable)



## **Parametrization of $\beta_t$**

Typically a sequence of linearly increasing constants: e.g. from  $\beta_1 = 10^{-4}$  to  $\beta_T = 0.02$

Can be fixed (not learnable)

## **Parametrization of reverse process variance $\Sigma_\theta$**

## Parametrization of $\beta_t$

Typically a sequence of linearly increasing constants: e.g. from  $\beta_1 = 10^{-4}$  to  $\beta_T = 0.02$

Can be fixed (not learnable)

## Parametrization of reverse process variance $\Sigma_\theta$

Can also be fixed  $\Sigma_\theta(x_t, t) = \sigma_t^2 \mathbf{I}$ , where  $\sigma_t$  is set to be a function of  $\beta_t$

## Parametrization of $\beta_t$

Typically a sequence of linearly increasing constants: e.g. from  $\beta_1 = 10^{-4}$  to  $\beta_T = 0.02$

Can be fixed (not learnable)

## Parametrization of reverse process variance $\Sigma_\theta$

Can also be fixed  $\Sigma_\theta(x_t, t) = \sigma_t^2 \mathbf{I}$ , where  $\sigma_t$  is set to be a function of  $\beta_t$

Alternative  $\Sigma_\theta(x_t, t) = \exp(v \log \beta_t + (1 - v) \log \tilde{\beta}_t)$ ,  $\tilde{\beta}_t = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}$ ,  $v$  is learnable

## Parametrization of $\beta_t$

Typically a sequence of linearly increasing constants: e.g. from  $\beta_1 = 10^{-4}$  to  $\beta_T = 0.02$

Can be fixed (not learnable)

## Parametrization of reverse process variance $\Sigma_\theta$

Can also be fixed  $\Sigma_\theta(x_t, t) = \sigma_t^2 \mathbf{I}$ , where  $\sigma_t$  is set to be a function of  $\beta_t$

Alternative  $\Sigma_\theta(x_t, t) = \exp(v \log \beta_t + (1 - v) \log \tilde{\beta}_t)$ ,  $\tilde{\beta}_t = \frac{1 - \bar{\alpha}_{t-1}}{1 - \bar{\alpha}_t}$ ,  $v$  is learnable

Hence, the loss  $L = L_{simple} + \lambda L_{VLB}$ ,  $\lambda$  is small  $\sim 0.001$  and  $L_{VLB}$  only guides the training of  $\Sigma_\theta$

(In  $L_{VLB}$  stop gradient with respect to  $\mu_\theta$ )

## Bonus! Score networks and guided diffusion

The score of each sample's  $x$  probability density function is defined as  $\nabla_x \log q(x)$

## Bonus! Score networks and guided diffusion

The score of each sample's  $x$  probability density function is defined as  $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score  $\nabla_x \log q(x)$  in an iterative process.

## Bonus! Score networks and guided diffusion

The score of each sample's  $x$  probability density function is defined as  $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score  $\nabla_x \log q(x)$  in an iterative process.

Score network  $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

## Bonus! Score networks and guided diffusion

The score of each sample's  $x$  probability density function is defined as  $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score  $\nabla_x \log q(x)$  in an iterative process.

Score network  $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

$$\text{If } x \sim \mathcal{N}(\mu, \sigma^2 \mathbf{I}), \quad \nabla_x \log p(x) = \nabla_x \left( -\frac{1}{2\sigma^2} (x - \mu)^2 \right) = -\frac{x - \mu}{\sigma^2} = -\frac{\varepsilon}{\sigma}, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I})$$



## Bonus! Score networks and guided diffusion

The score of each sample's  $x$  probability density function is defined as  $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score  $\nabla_x \log q(x)$  in an iterative process.

Score network  $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

$$\text{If } x \sim \mathcal{N}(\mu, \sigma^2 \mathbf{I}), \quad \nabla_x \log p(x) = \nabla_x \left( -\frac{1}{2\sigma^2} (x - \mu)^2 \right) = -\frac{x - \mu}{\sigma^2} = -\frac{\varepsilon}{\sigma}, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I})$$

$$q(x_t | x_0) \sim \mathcal{N}(\sqrt{\bar{\alpha}_t} x_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

## Bonus! Score networks and guided diffusion

The score of each sample's  $x$  probability density function is defined as  $\nabla_x \log q(x)$

Langevin dynamics can sample data points from a probability density distribution using only the score  $\nabla_x \log q(x)$  in an iterative process.

Score network  $s_\theta(x_t, t) \approx \nabla_x \log q(x)$

$$\text{If } x \sim \mathcal{N}(\mu, \sigma^2 \mathbf{I}), \quad \nabla_x \log p(x) = \nabla_x \left( -\frac{1}{2\sigma^2} (x - \mu)^2 \right) = -\frac{x - \mu}{\sigma^2} = -\frac{\varepsilon}{\sigma}, \quad \varepsilon \sim \mathcal{N}(0, \mathbf{I})$$

$$q(x_t | x_0) \sim \mathcal{N}(\sqrt{\bar{\alpha}_t} x_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

$$\text{And therefore } s_\theta(x_t, t) \approx \nabla_{x_t} \log q(x_t) = E_{q(x_0)} \nabla_{x_t} q(x_t | x_0) = E_{q(x_0)} \left[ -\frac{\varepsilon_\theta(x_t, t)}{\sqrt{1 - \bar{\alpha}_t}} \right] = -\frac{\varepsilon_\theta(x_t, t)}{\sqrt{1 - \bar{\alpha}_t}}$$

# Guided diffusion

- We have additional input  $y$  (a class label in classifier guided diffusion)
- We want to model a conditional distribution  $p(x | y)$  instead



Conditioned on dogs

## Classifier guided diffusion

We separately train a classifier  $f_\phi(y \mid x_t, t)$  on a noisy image  $x_t$ , and use gradients  $\nabla_x \log f_\phi(y \mid x_t)$  to guide the diffusion. Let us have a joint distribution  $q(x_t, y)$ ,  $y$  is e.g. the image label

## Classifier guided diffusion

We separately train a classifier  $f_\phi(y \mid x_t, t)$  on a noisy image  $x_t$ , and use gradients  $\nabla_x \log f_\phi(y \mid x_t)$

to guide the diffusion. Let us have a joint distribution  $q(x_t, y)$ ,  $y$  is e.g. the image label

$$\nabla_{x_t} \log q(x_t \mid y) = \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y \mid x_t) - \nabla_{x_t} \log q(y)$$

## Classifier guided diffusion

We separately train a classifier  $f_\phi(y | x_t, t)$  on a noisy image  $x_t$ , and use gradients  $\nabla_x \log f_\phi(y | x_t)$

to guide the diffusion. Let us have a joint distribution  $q(x_t, y)$ ,  $y$  is e.g. the image label

$$\nabla_{x_t} \log q(x_t | y) = \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y | x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y | x_t)$$

## Classifier guided diffusion

We separately train a classifier  $f_\phi(y \mid x_t, t)$  on a noisy image  $x_t$ , and use gradients  $\nabla_x \log f_\phi(y \mid x_t)$  to guide the diffusion. Let us have a joint distribution  $q(x_t, y)$ ,  $y$  is e.g. the image label

$$\begin{aligned}\nabla_{x_t} \log q(x_t \mid y) &= \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y \mid x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y \mid x_t) \\ &= -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} \nabla_{x_t} \log f_\phi(y \mid x_t))\end{aligned}$$

## Classifier guided diffusion

We separately train a classifier  $f_\phi(y | x_t, t)$  on a noisy image  $x_t$ , and use gradients  $\nabla_x \log f_\phi(y | x_t)$  to guide the diffusion. Let us have a joint distribution  $q(x_t, y)$ ,  $y$  is e.g. the image label

$$\begin{aligned}\nabla_{x_t} \log q(x_t | y) &= \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y | x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y | x_t) \\ &= -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} \nabla_{x_t} \log f_\phi(y | x_t))\end{aligned}$$

Classifier guided predictor  $\bar{\varepsilon}_\theta(x_t, t) = \varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} w \nabla_{x_t} \log f_\phi(y | x_t)$



## Classifier guided diffusion

We separately train a classifier  $f_\phi(y | x_t, t)$  on a noisy image  $x_t$ , and use gradients  $\nabla_x \log f_\phi(y | x_t)$  to guide the diffusion. Let us have a joint distribution  $q(x_t, y)$ ,  $y$  is e.g. the image label

$$\begin{aligned}\nabla_{x_t} \log q(x_t | y) &= \nabla_{x_t} \log q(x_t) + \nabla_{x_t} \log q(y | x_t) - \nabla_{x_t} \log q(y) \approx -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} \varepsilon_\theta(x_t, t) + \nabla_{x_t} \log f_\phi(y | x_t) \\ &= -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} \nabla_{x_t} \log f_\phi(y | x_t))\end{aligned}$$

Classifier guided predictor  $\bar{\varepsilon}_\theta(x_t, t) = \varepsilon_\theta(x_t, t) - \sqrt{1 - \bar{\alpha}_t} w \nabla_{x_t} \log f_\phi(y | x_t)$

Where  $w$  is the strength of the guidance

# Classifier guided diffusion

---

**Algorithm 1** Classifier guided diffusion sampling, given a diffusion model  $(\mu_\theta(x_t), \Sigma_\theta(x_t))$ , classifier  $f_\phi(y|x_t)$ , and gradient scale  $s$ .

---

Input: class label  $y$ , gradient scale  $s$

$x_T \leftarrow$  sample from  $\mathcal{N}(0, \mathbf{I})$

**for all**  $t$  from  $T$  to 1 **do**

$\mu, \Sigma \leftarrow \mu_\theta(x_t), \Sigma_\theta(x_t)$

$x_{t-1} \leftarrow$  sample from  $\mathcal{N}(\mu + s\Sigma \nabla_{x_t} \log f_\phi(y|x_t), \Sigma)$

**end for**

**return**  $x_0$

---

Downsides:

# Classifier guided diffusion

---

**Algorithm 1** Classifier guided diffusion sampling, given a diffusion model  $(\mu_\theta(x_t), \Sigma_\theta(x_t))$ , classifier  $f_\phi(y|x_t)$ , and gradient scale  $s$ .

---

Input: class label  $y$ , gradient scale  $s$

$x_T \leftarrow$  sample from  $\mathcal{N}(0, \mathbf{I})$

**for all**  $t$  from  $T$  to 1 **do**

$\mu, \Sigma \leftarrow \mu_\theta(x_t), \Sigma_\theta(x_t)$

$x_{t-1} \leftarrow$  sample from  $\mathcal{N}(\mu + s\Sigma \nabla_{x_t} \log f_\phi(y|x_t), \Sigma)$

**end for**

**return**  $x_0$

---

Downsides:

- The classifier has to cope with noise, might need to be trained separately

# Classifier guided diffusion

---

**Algorithm 1** Classifier guided diffusion sampling, given a diffusion model  $(\mu_\theta(x_t), \Sigma_\theta(x_t))$ , classifier  $f_\phi(y|x_t)$ , and gradient scale  $s$ .

---

Input: class label  $y$ , gradient scale  $s$

$x_T \leftarrow$  sample from  $\mathcal{N}(0, \mathbf{I})$

**for all**  $t$  from  $T$  to 1 **do**

$\mu, \Sigma \leftarrow \mu_\theta(x_t), \Sigma_\theta(x_t)$

$x_{t-1} \leftarrow$  sample from  $\mathcal{N}(\mu + s\Sigma \nabla_{x_t} \log f_\phi(y|x_t), \Sigma)$

**end for**

**return**  $x_0$

---

Downsides:

- The classifier has to cope with noise, might need to be trained separately
- If noise-robust might be inefficient: most of the information on  $x$  is irrelevant to  $y$

## Classifier free guidance

- Train a diffusion model  $p(x | y)$  with conditioning dropout: conditional information is removed some percentage of the time

## Classifier free guidance

- Train a diffusion model  $p(x | y)$  with conditioning dropout: conditional information is removed some percentage of the time
- Resulting model can be conditional  $p(x | y)$  and unconditional  $p(x)$

## Classifier free guidance

- Train a diffusion model  $p(x | y)$  with conditioning dropout: conditional information is removed some percentage of the time
- Resulting model can be conditional  $p(x | y)$  and unconditional  $p(x)$
- No need for a separate classifier!

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$



# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$
- Conditional model  $p_\theta(x | y)$  parametrised through  $\varepsilon_\theta(x_t, t, y)$

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$
- Conditional model  $p_\theta(x | y)$  parametrised through  $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data  $(x, y)$

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$
- Conditional model  $p_\theta(x | y)$  parametrised through  $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data  $(x, y)$
- Conditional information  $y$  gets discarded periodically:  $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$
- Conditional model  $p_\theta(x | y)$  parametrised through  $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data  $(x, y)$
- Conditional information  $y$  gets discarded periodically:  $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t)$$

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$
- Conditional model  $p_\theta(x | y)$  parametrised through  $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data  $(x, y)$
- Conditional information  $y$  gets discarded periodically:  $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t, y) - \varepsilon_\theta(x_t, t))$$

# Classifier free guidance

Guide the diffusion without an independent classifier  $f_\phi$

- Unconditional denoising diffusion model  $p_\theta(x)$ , parametrised through a score estimator  $\varepsilon_\theta(x_t, t)$
- Conditional model  $p_\theta(x_t | y)$  parametrised through  $\varepsilon_\theta(x_t, t, y)$
- Conditional model is trained on paired data  $(x, y)$
- Conditional information  $y$  gets discarded periodically:  $\varepsilon_\theta(x_t, t) = \varepsilon_\theta(x_t, t, y = \emptyset)$

$$\nabla_{x_t} \log p(y | x_t) = \nabla_{x_t} \log p(x_t | y) - \nabla_{x_t} \log p(x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_\theta(x_t, t, y) - \varepsilon_\theta(x_t, t))$$

$$\text{Now } \nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t) + \nabla_{x_t} \log p(y | x_t) = \nabla_x \log p(x_t) - \frac{1}{\sqrt{1 - \bar{\alpha}}} (\varepsilon_\theta(x_t, t, y) - \varepsilon_\theta(x_t, t))$$

## Classifier free guidance

$$\nabla_{x_t} \log p(x_t, y) = \nabla_x \log p(x_t) - \frac{1}{\sqrt{1 - \bar{\alpha}}} (\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t)) =$$

$$\varepsilon_{\theta}(x, t) - w(\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t)) = (1 + w)\varepsilon_{\theta}(x, t) - w\varepsilon_{\theta}(x_t, t, y)$$



## Classifier free guidance

$$\nabla_{x_t} \log p(x_t, y) = \nabla_{x_t} \log p(x_t | y) + \nabla_{x_t} \log p(y | x_t)$$

$$\nabla_{x_t} \log p(y | x_t) = -\frac{1}{\sqrt{1 - \bar{\alpha}_t}} (\varepsilon_{\theta}(x_t, t, y) - \varepsilon_{\theta}(x_t, t))$$

Hence, analogously to the case of classifier guided diffusion